



PATIENT FACT SHEET

Teriparatide (Forteo)



WHAT IS IT?


Teriparatide is used in post-menopausal women and men who have osteoporosis and are at high risk for bone fractures. It is often used when other treatments fail or in severe forms of the disease. Teriparatide itself builds bone and is different from other forms of osteoporosis treatment, which prevent bone breakdown.



HOW TO TAKE IT

Teriparatide is given by injection daily for two years. The medication comes in an injector, which is like a pen. To inject the medicine, place the injector on the thigh or upper arm, push the button and it is automatically

injected. The dose is 20mcg a day and this is preset by the pen. The needle size is the same as an insulin needle. The injector has enough medicine for 28 days, and the medication should be kept refrigerated.



SIDE EFFECTS

Patients may get dizzy within four hours after taking the first few doses, but this goes away within a few hours. Occasionally, the injection site can become itchy, red, or swollen. Teriparatide rarely can cause increased levels of calcium in the blood, nausea, or achy joints.

Teriparatide has a black box warning, meaning the Food and Drug Administration is concerned with an increased risk of cancer. Sarcoma, a type of cancer, was found in rats that were given very large doses of teriparatide. Repeat studies in monkeys did not reveal this and since it has been released in humans, there has only been one case of sarcoma in a patient taking teriparatide.



TELL YOUR DOCTOR

You should notify your doctor if you have trouble injecting yourself; if the injection site becomes red, itchy, warm, or swollen; or if you have increasing achy joints or muscle spasms.

Make sure to notify your other physicians while you are taking this drug. If you are pregnant or considering

pregnancy, let your doctor know before starting this medication. Women should discuss birth control with their primary care physicians or gynecologists. Breastfeeding should be avoided while taking teriparatide because the drug can enter breast milk.

Be sure to talk with your doctor before undergoing any surgeries while taking this medication.